

Original Paper

Using Implementation Science to Understand Tele dermatology Implementation Early in the COVID-19 Pandemic: Cross-sectional Study

Shanelle Mariah Briggs¹, BS, BA; Jules Benjamin Lipoff², MD; Sigrid Marie Collier³, MPH, MD

¹School of Medicine, University of Washington, Seattle, WA, United States

²Department of Dermatology, University of Pennsylvania, Philadelphia, PA, United States

³Division of Dermatology, University of Washington, Seattle, WA, United States

Corresponding Author:

Sigrid Marie Collier, MPH, MD

Division of Dermatology

University of Washington

BB-1353 Health Sciences Center, Box 356524

1959 NE Pacific Street

Seattle, WA, 98195-6524

United States

Phone: 1 206 543 5290

Fax: 1 302 202 3331

Email: coll0640@uw.edu

Abstract

Background: Implementation science has been recognized for its potential to improve the integration of evidence-based practices into routine dermatologic care. The COVID-19 pandemic has resulted in rapid tele dermatology implementation worldwide. Although several studies have highlighted patient and care provider satisfaction with tele dermatology during the COVID-19 pandemic, less is known about the implementation process.

Objective: Our goal was to use validated tools from implementation science to develop a deeper understanding of the implementation of tele dermatology during the COVID-19 pandemic. Our primary aims were to describe (1) the acceptability and feasibility of the implementation of tele dermatology and (2) organizational readiness for the implementation of tele dermatology during the COVID-19 pandemic. We also sought to offer an example of how implementation science can be used in dermatologic research.

Methods: An anonymous, web-based survey was distributed to Association of Professors of Dermatology members. It focused on (1) the acceptability, feasibility, and appropriateness of tele dermatology and (2) organizational readiness for implementing tele dermatology. It incorporated subscales from the Organizational Readiness to Change Assessment—a validated measure of organizational characteristics that predict implementation success.

Results: Of the 518 dermatologists emailed, 35 (7%) responded, and all implemented or scaled up tele dermatology during the pandemic. Of the 11 care providers with the highest level of organizational readiness, 11 (100%) said that they plan to continue using tele dermatology after the pandemic. Most respondents agreed or strongly agreed that they had sufficient training (24/35, 69%), financial resources (20/35, 57%), and facilities (20/35, 57%). However, of the 35 respondents, only 15 (43%) agreed or strongly agreed that they had adequate staffing support. Most respondents considered the most acceptable tele dermatology modality to be synchronous audio and video visits with supplemental stored digital photos (23/35, 66%) and considered the least acceptable modality to be telephone visits without stored digital photos (6/35, 17%). Overall, most respondents thought that the implementation of synchronous audio and video with stored digital photos (31/35, 89%) and telephone visits with stored digital photos (31/35, 89%) were the most feasible. When asked about types of visits that were acceptable for synchronous video/audio visits (with stored digital photos), 18 of the 31 respondents (58%) said “new patients,” 27 (87%) said “existing patients,” 19 (61%) said “medication monitoring,” 3 (10%) said “total body skin exams,” and 22 (71%) said “lesions of concern.”

Conclusions: This study serves as an introduction to how implementation science research methods can be used to understand the implementation of novel technologies in dermatology. Our work builds upon prior studies by further characterizing the

acceptability and feasibility of different teledermatology modalities. Our study may suggest initial insights on how dermatology practices and health care systems can support dermatologists in successfully incorporating teledermatology after the pandemic.

(*JMIR Dermatol* 2022;5(2):e33833) doi: [10.2196/33833](https://doi.org/10.2196/33833)

KEYWORDS

teledermatology; telemedicine; telehealth; COVID-19; remote care; implementation science; store and forward; Organizational Readiness to Change Assessment; acceptability; feasibility; digital health; dermatology; dermatologist; health technology

Introduction

When the COVID-19 pandemic began in the spring of 2020, restrictions were placed on in-person visits. This crisis led to the rapid increase in teledermatology implementation, which was made possible by policy changes that overcame prior barriers to implementation, such as the lack of insurance reimbursement, liability concerns, and licensing restrictions [1,2]. A recent review highlighted teledermatology's potential to reduce health care disparities in underserved and marginalized communities, calling for future efforts to study implementation, as teledermatology has expanded greatly during the pandemic [3]. Although much of the published work on teledermatology during the COVID-19 pandemic focused on satisfaction among patients and care providers, less is known about its actual implementation during the early months of the pandemic [4,5]. The field of implementation science has been recognized for its potential to improve the integration of evidence-based practices into routine dermatologic care [6].

In order for teledermatology to be successfully incorporated into routine dermatologic practice, there must be buy-in from dermatology patients, dermatologists, and health systems. Our study adds to the growing body of evidence for teledermatology by using validated implementation science tools to develop a deeper understanding of the implementation of teledermatology during the COVID-19 pandemic.

Implementation science uses specific terminology to describe key predictors of and outcomes for the implementation of evidence-based interventions. Implementation outcomes refer to “the effects of deliberate and purposive actions to implement new treatments, practices, and services,” and they “[serve] as an indicator of implementation success” [7]. In this study, we focused on evaluating the *acceptability* and *feasibility* of teledermatology, which are defined, respectively, as follows: (1) “the perception among implementation stakeholders that a given treatment, service, practice, or innovation is agreeable, palatable, or satisfactory” and (2) “the extent to which a new treatment, or innovation, can be successfully used or carried out within a given agency or setting” [7]. We also evaluated *organizational readiness* for change, which is defined as “the extent to which organizational members are psychologically and behaviourally prepared to implement organizational change” [8].

We use the lens of implementation science to describe teledermatology implementation. The objective of our study

was to understand the acceptability and feasibility of the implementation of teledermatology during the COVID-19 pandemic, as well as organizational readiness for the implementation of teledermatology. We also sought to offer an example of how implementation science can be used in dermatologic research.

Methods

Ethical Considerations

Our study was deemed exempt from ethics approval by the University of Washington Human Subjects Division and the University of Washington Institutional Review Board (IRB ID: STUDY00010266).

Study Design

We conducted a cross-sectional study of dermatologists' perceptions of teledermatology implementation during the COVID-19 pandemic. We surveyed members of the Association of Professors of Dermatology (APD) between November 20 and December 9, 2020 ([Multimedia Appendix 1](#)). An initial email was sent on November 20, and it was resent on December 2 to try to increase the response rate. The survey focused on (1) the acceptability, feasibility, and appropriateness of teledermatology and (2) organizational readiness for implementing teledermatology. We used an abbreviated, single-item version of a validated scale [9] to assess the acceptability of different teledermatology modalities (eg, synchronous audio and video and stored digital photos). Using the same scale, we assessed the acceptability of teledermatology for different dermatologic conditions and purposes (eg, lesions of concern and medication monitoring). We also incorporated selected subscales from the validated Organizational Readiness to Change Assessment (ORCA) instrument (eg, culture and resources). There were 26 ORCA questions, which were scored on a 5-point Likert scale ranging from “Strongly Disagree” to “Strongly Agree.” The possible ORCA scores ranged from 26 to 130, with higher scores indicating higher organizational readiness for change. The entire survey was tested for face validity and readability through pilots with dermatologists. It was iteratively refined based on their feedback. The survey was administered via email, and responses were collected anonymously. ORCA scores were reported as unweighted composite scores, and participants were stratified by ORCA score tertiles—“low,” “medium,” and “high” organizational readiness for change ([Table 1](#)). We did not perform statistical hypothesis testing, in accordance with best practices [10].

Table 1. Demographics and outcomes by Organizational Readiness to Change Assessment (ORCA) tertiles.

	ORCA score tertiles			Total (N=35)
	Low (n=11)	Medium (n=13)	High (n=11)	
Sex, n (%)				
Male	2 (18)	6 (47)	5 (46)	13 (37)
Female	9 (82)	7 (54)	6 (55)	22 (63)
Race, n (%)				
American Indian or Alaska Native	1 (9)	0 (0)	0 (0)	1 (3)
Asian	1 (9)	2 (15)	2 (18)	5 (14)
Black or African American	1 (9)	0 (0)	0 (0)	1 (3)
Hispanic, Latinx, or Spanish origin	1 (9)	2 (15)	1 (9)	4 (11)
White	7 (64)	9 (69)	8 (73)	24 (69)
Practice, n (%)				
Dermatology group practice	0 (0)	0 (0)	2 (18)	2 (6)
Multispecialty group practice	1 (9)	0 (0)	0 (0)	1 (3)
Academic practice	9 (82)	12 (92)	9 (82)	30 (86)
Veterans administration and academic practice	1 (9)	1 (8)	0 (0)	2 (6)
Number of years in practice				
Years, mean (SD)	3.91 (1.97)	3.69 (1.75)	3.82 (1.78)	3.80 (1.78)
Years, median (quartile 1, quartile 3)	4.00 (2.50, 6.00)	3.00 (2.00, 6.00)	4.00 (3.00, 5.00)	4.00 (2.00, 6.00)
Total ORCA score				
Score, mean (SD)	70.0 (9.58)	90.4 (5.36)	110.7 (8.75)	90.4 (18.1)
Score, median (quartile 1, quartile 3)	68.00 (66.0, 78.0)	91.0 (88.0, 96.0)	110.0 (104.0, 113.5)	91.0 (78.5, 102.5)
Stored digital photos alone are acceptable, n (%)				
Completely agree or agree	7 (64)	5 (39)	6 (55)	18 (51)
Completely disagree, disagree, or neither agree nor disagree	4 (36)	8 (62)	5 (46)	17 (49)
Telephone visits without photos are acceptable, n (%)				
Completely agree or agree	2 (18)	1 (8)	3 (27)	6 (17)
Completely disagree, disagree, or neither agree nor disagree	9 (82)	12 (92)	8 (73)	29 (83)
Telephone visits with photos are acceptable, n (%)				
Completely agree or agree	7 (64)	7 (54)	6 (55)	20 (57)
Completely disagree, disagree, or neither agree nor disagree	4 (36)	6 (46)	5 (46)	15 (43)
Synchronous audio and video visits without photos are acceptable, n (%)				
Completely agree or agree	3 (27)	4 (31)	5 (46)	12 (34)
Completely disagree, disagree, or neither agree nor disagree	8 (73)	9 (69)	6 (55)	23 (66)
Synchronous audio and video visits with photos are acceptable, n (%)				
Completely agree or agree	4 (36)	11 (85)	8 (73)	23 (66)
Completely disagree, disagree, or neither agree nor disagree	7 (64)	2 (15)	3 (27)	12 (34)
Plan to use telemedicine after the pandemic, n (%)				
Yes	10 (91)	12 (92)	11 (100)	33 (94)
No	1 (9)	1 (8)	0 (0)	2 (6)

Results

Of the 518 dermatologists on the APD email listserv, 35 (7%) responded, and all implemented or scaled up teledermatology during the pandemic. Of the 35 respondents, 35 (100%) said that the peak use of teledermatology occurred between the months of April and September 2020. Thus, all respondents had completed the initial implementation by the time of survey distribution in December 2020. Further, 94% (33/35) plan to continue using teledermatology after the pandemic. The benefits of teledermatology included less travel time and expense for patients (n=35, 100%), continued patient care (n=33, 94%), the ability to avoid the risk of infection (n=35, 100%), and work flexibility (n=27, 77%). Respondents also experienced challenges with teledermatology, including technology issues (n=22, 63%) and challenges with caring for older adults (n=18, 51%). All 11 care providers with “high” ORCA scores said that they plan to continue using teledermatology after the pandemic (Table 1). With regard to organizational readiness for the implementation of teledermatology during the pandemic, 24 of the 35 care providers (69%) agreed or strongly agreed that they had sufficient training, 20 (57%) had sufficient financial resources, and 20 (57%) had sufficient facilities. Most respondents had care provider buy-in (25/35, 71%) and felt that teledermatology implementation took into consideration the needs and preferences of patients (27/35, 77%). On the other hand, fewer respondents had a dedicated team for implementing the intervention (14/34, 41%), had sufficient staffing support (15/35, 43%), or had successfully piloted telemedicine prior to the pandemic (13/35, 37%). Most of the 35 respondents reported using several implementation strategies, which included a dedicated clinical champion (n=26, 74%); feedback to clinicians (n=20, 59%); education (n=24, 69%); and, less commonly, staff incentives (n=4, 11%). Table S1 in [Multimedia Appendix 2](#) shows these supplemental results. Of the 11 respondents with “low” ORCA scores, 6 (55%) agreed or strongly agreed with the ORCA components about having a clinical champion, 5 (45%) agreed or strongly agreed with giving feedback to clinicians, 2 (18%) agreed or strongly agreed with education, and 0 (0%) agreed or strongly agreed with staff incentives. Of the 11 participants with “high” ORCA scores, 11 (100%) agreed or strongly agreed with having a clinical champion, 9 (82%) agreed or strongly agreed with giving feedback to clinicians, 10 (91%) agreed or strongly agreed with education, and 4 (36%) agreed or strongly agreed with staff incentives (Table S2 in [Multimedia Appendix 2](#)).

The most acceptable teledermatology modality was synchronous audio and video visits with stored digital photos (23/35, 66%). The least acceptable modality was telephone visits without stored digital photos (6/35, 17%). When comparing participants with “low” ORCA scores to those with “medium” and “high” ORCA scores, synchronous audio and video visits with stored digital photos were less acceptable among those with “low” ORCA scores (4/11, 36%) relative to those with “medium” (11/13, 85%) and “high” (8/11, 73%) ORCA scores. However, the acceptability of consultations involving stored digital photos was higher among those with “low” ORCA scores (7/11, 64%) relative to those with “medium” (5/13, 39%) and “high” (6/11,

55%) ORCA scores. Along with acceptability (Table 1), feasibility was also addressed. Overall, among the 35 respondents, synchronous audio and video visits with stored digital photos (n=31, 89%) and telephone visits with stored digital photos (n=31, 89%) were deemed the most feasible teledermatology modalities. Other modalities were also deemed feasible, though less so, including consultations involving stored digital photos (n=26, 74%), synchronous audio and video visits without stored digital photos (n=23, 66%), and telephone visits without stored digital photos (n=21, 60%). When asked about types of visits that were acceptable for synchronous video/audio visits (with stored digital photos), 18 of the 31 respondents (58%) said “new patients,” 27 (87%) said “existing patients,” 19 (61%) said “medication monitoring,” 3 (10%) said “total body skin exams,” and 22 (71%) said “lesions of concern. The majority of surveyed dermatologists felt that synchronous video/audio (without stored digital photos) was acceptable for “existing patients” (29/32, 91%) and “medication monitoring” (29/32, 91%). Fewer respondents felt that synchronous video/audio (without stored digital photos) was acceptable for “new patients” (12/32, 38%), and very few felt that teledermatology was acceptable for “lesions of concern” (5/32, 16%) and “total body skin exams” (2/32, 6%). Additional detailed results are shown in [Table 1](#).

Discussion

Principal Results

In our study, although most dermatologists (33/35, 94%) planned to continue using teledermatology after the pandemic, there was some indication that they lacked support in certain areas (eg, staffing and facilities) during implementation early in the COVID-19 pandemic. Additionally, not all teledermatology modalities were equally acceptable or feasible. Among respondents, telephone and synchronous audio and video visits were the least acceptable and feasible modalities, whereas modalities that combined stored digital photos with telephone visits or synchronous audio and video visits were the most acceptable and feasible modalities.

Teledermatology has been a part of dermatologic care for over 25 years [11]. Although consultations involving stored digital photos (store and forward) have historically been the dominant teledermatology modality in clinical practice [12], the COVID-19 pandemic has resulted in the rapid implementation of synchronous audio and video teledermatology [13].

Although the majority of dermatologists (33/35, 94%) in this study planned to continue using teledermatology after the pandemic, there is likely some variability in intentions to continue using teledermatology, depending on the population surveyed and timing. An earlier survey by the American Academy of Dermatology in May 2020, which included a larger proportion of private practice dermatologists, found that just over half (58%) of dermatologists planned to continue using teledermatology after the pandemic [13]. Despite the differences, both surveys highlight the importance of teledermatology in the future, with over 50% of dermatologists intending to practice teledermatology.

Overall, respondents supported the use of tele dermatology after the pandemic; however, we found that both telephone visits and synchronous audio and video visits without stored digital photos were the two least acceptable and feasible modalities. This finding aligns with patients' experiences with tele dermatology. Despite high levels of patient satisfaction and willingness to continue using tele dermatology after the pandemic [4,5], satisfaction with dermatology telephone visits is lower, and fewer dermatology patients are willing to use telephone visits for future dermatologic care [14]. Taken as a whole, our findings build on a growing body of evidence that certain modalities, particularly telephone visits, are less acceptable to both patients and care providers. Dermatologists rely upon the clear and accurate visualization of the skin, which telephone and synchronous video/audio visits alone may not offer.

Incorporating stored digital photos may overcome some of the limitations of using synchronous audio and video visits and telephone visits in isolation. Acceptability was higher for including stored digital photos with synchronous audio and video visits or telephone visits when compared to that for visits without stored digital photos and consultations involving stored digital photos alone. Prior studies that were conducted during the COVID-19 pandemic showed that care providers were split between preferring synchronous (54%) and asynchronous (46%) modalities but did not assess the combination of asynchronous and synchronous approaches [15]. Dermatologists may be able to maximize the benefits of synchronous modalities (synchronous audio and video visits and telephone visits) and asynchronous modalities (stored digital photos) by combining them to create a more acceptable and preferable tele dermatology experience.

We also found that there may be variability in the acceptability of tele dermatology based on the type of dermatologic condition and visit. The majority of surveyed dermatologists felt that synchronous video/audio (without stored digital photos) was acceptable for "existing patients" (29/32, 91%) and "medication monitoring" (29/32, 91%). Fewer respondents felt that synchronous video/audio (without stored digital photos) was acceptable for "new patients" (12/32, 38%), and very few felt that tele dermatology was acceptable for "lesions of concern" (5/32, 16%) and "total body skin exams" (2/32, 6%). The American Academy of Dermatology's survey found a similarly low number of dermatologists who were comfortable with performing total body skin exams via tele dermatology, with 96% believing that this requires an in-person examination [13]. This work adds to our understanding of the types of patient concerns for which tele dermatology is the most acceptable. We hope that tele dermatology guidelines for best practices can evolve via this growing collection of work.

When implementing novel health care technologies (including tele dermatology technologies), organizational factors, such as organizational readiness for change, are important determinants of implementation success [16]. We found that although support for tele dermatology implementation was high in most areas (eg, training and care provider buy-in), dermatologists lacked

organizational support in other areas, such as staffing support and facilities. In addition, we found that respondents with a "low" organizational readiness for change tended to find synchronous audio and video visits less acceptable when compared to respondents with "medium" and "high" ORCA scores. It is plausible that in dermatology practices with a lower organizational readiness for change, limited support for newly implementing synchronous audio and video tele dermatology [13] resulted in negative experiences during the COVID-19 pandemic. This hints at the potential importance of organizational readiness in determining the success of tele dermatology implementation. Although specific implementation process details were outside the scope of this work, lower scores for important implementation strategies, including having a clinical champion, giving feedback to clinicians, and providing education, contributed to respondents having "low" scores for organizational readiness for change, and this may provide clues as to the specific implementation strategies that are important for the successful implementation of synchronous audio and video tele dermatology early in the COVID-19 pandemic. Future research will be needed to explore the roles of specific implementation strategies, implementation processes, and costs in determining the success of newly implemented tele dermatology programs.

We acknowledge that this cross-sectional survey has significant limitations, given its modest sample size and response rate. Therefore, we cannot draw definitive conclusions on the associations between organizational readiness for change and tele dermatology implementation outcomes. Still, our total of 35 respondents and response rate of 7% (35/518) are similar to those of other nonincentivized physician surveys [13]. For these reasons, the findings of this survey may not be generalizable. As the survey was distributed to the APD, the majority of respondents (32/35) practiced in academic dermatology settings. Thus, respondents with an interest in tele dermatology may have been overrepresented. Additionally, respondents were mostly White (24/35, 69%); as such, the opinions of dermatologists from all backgrounds were not captured. Despite these limitations, this work provides valuable descriptive insights into the role of implementation science in understanding tele dermatology implementation during the COVID-19 pandemic.

Conclusions

This study serves as an introduction to how implementation science research methods can be used to understand the implementation of novel technologies in dermatology. Our work builds on prior work by further characterizing the acceptability and feasibility of different tele dermatology modalities. Our study also contributes initial insights on how dermatology practices and health care systems can support dermatologists in successfully incorporating tele dermatology after the pandemic. Finally, this work highlights newer methods for identifying organizational factors that can be optimized to improve future tele dermatology implementation efforts.

Acknowledgments

The research reported in this publication was supported by the Fogarty International Center and National Cancer Institute of the National Institutes of Health under grant D43TW009345-09S7, which was awarded to the Northern Pacific Global Health Fellows Program. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. We give special thanks to Dr Michi Shinohara, Dr Cuong Nguyen, Dr John Olerud, Dr Markus Boos, Dr Jay Vary, Dr Nicholas Compton, Dr April Schachtel, and Dr Max Vale for their contributions.

SMC's time on this project was funded by the National Institutes of Health under grant D43TW009345-09S7.

Conflicts of Interest

None declared.

Multimedia Appendix 1

Survey instrument and recruitment letter/email.

[\[DOCX File , 5610 KB-Multimedia Appendix 1\]](#)

Multimedia Appendix 2

Supplemental results.

[\[DOCX File , 32 KB-Multimedia Appendix 2\]](#)

References

1. Yeboah CB, Harvey N, Krishnan R, Lipoff JB. The impact of COVID-19 on tele dermatology: A review. *Dermatol Clin* 2021 Oct;39(4):599-608 [FREE Full text] [doi: [10.1016/j.det.2021.05.007](https://doi.org/10.1016/j.det.2021.05.007)] [Medline: [34556249](https://pubmed.ncbi.nlm.nih.gov/34556249/)]
2. Ramírez-Cornejo C, Muñoz-López C, Del Barrio-Díaz P, Jaque A, Majerson D, Navarrete-Dechent C, Grupo de Tele dermatología UC-Christus. [Rapid implementation of tele-dermatology during COVID-19 pandemic in an academic dermatology department]. *Rev Med Chil* 2021 Oct;149(10):1467-1472 [FREE Full text] [doi: [10.4067/s0034-98872021001001467](https://doi.org/10.4067/s0034-98872021001001467)] [Medline: [35319636](https://pubmed.ncbi.nlm.nih.gov/35319636/)]
3. Maddukuri S, Patel J, Lipoff JB. Tele dermatology addressing disparities in health care access: a review. *Curr Dermatol Rep* 2021 Mar 12:1-8 [FREE Full text] [doi: [10.1007/s13671-021-00329-2](https://doi.org/10.1007/s13671-021-00329-2)] [Medline: [33747638](https://pubmed.ncbi.nlm.nih.gov/33747638/)]
4. Turan Ç, Utlü Z. Investigating the factors enabling the accurate implementation of the patient-assisted tele dermatology model during the pandemic in Turkey: A pilot study. *Dermatol Ther* 2021 Mar;34(2):e14741. [doi: [10.1111/dth.14741](https://doi.org/10.1111/dth.14741)] [Medline: [33404147](https://pubmed.ncbi.nlm.nih.gov/33404147/)]
5. Hamad J, Fox A, Kammire MS, Hollis AN, Khairat S. Evaluating the experiences of new and existing tele dermatology patients during the COVID-19 pandemic: Cross-sectional survey study. *JMIR Dermatol* 2021 May 05;4(1):e25999 [FREE Full text] [doi: [10.2196/25999](https://doi.org/10.2196/25999)] [Medline: [34028471](https://pubmed.ncbi.nlm.nih.gov/34028471/)]
6. Ashrafzadeh S, Metlay JP, Choudhry NK, Emmons KM, Asgari MM. Using implementation science to optimize the uptake of evidence-based medicine into dermatology practice. *J Invest Dermatol* 2020 May;140(5):952-958 [FREE Full text] [doi: [10.1016/j.jid.2019.10.011](https://doi.org/10.1016/j.jid.2019.10.011)] [Medline: [31862108](https://pubmed.ncbi.nlm.nih.gov/31862108/)]
7. Proctor E, Silmere H, Raghavan R, Hovmand P, Aarons G, Bunger A, et al. Outcomes for implementation research: conceptual distinctions, measurement challenges, and research agenda. *Adm Policy Ment Health* 2011 Mar;38(2):65-76 [FREE Full text] [doi: [10.1007/s10488-010-0319-7](https://doi.org/10.1007/s10488-010-0319-7)] [Medline: [20957426](https://pubmed.ncbi.nlm.nih.gov/20957426/)]
8. Weiner BJ, Amick H, Lee SYD. Conceptualization and measurement of organizational readiness for change: a review of the literature in health services research and other fields. *Med Care Res Rev* 2008 Aug;65(4):379-436. [doi: [10.1177/1077558708317802](https://doi.org/10.1177/1077558708317802)] [Medline: [18511812](https://pubmed.ncbi.nlm.nih.gov/18511812/)]
9. Weiner BJ, Lewis CC, Stanick C, Powell BJ, Dorsey CN, Clary AS, et al. Psychometric assessment of three newly developed implementation outcome measures. *Implement Sci* 2017 Aug 29;12(1):108 [FREE Full text] [doi: [10.1186/s13012-017-0635-3](https://doi.org/10.1186/s13012-017-0635-3)] [Medline: [28851459](https://pubmed.ncbi.nlm.nih.gov/28851459/)]
10. Leon AC, Davis LL, Kraemer HC. The role and interpretation of pilot studies in clinical research. *J Psychiatr Res* 2011 May;45(5):626-629 [FREE Full text] [doi: [10.1016/j.jpsychires.2010.10.008](https://doi.org/10.1016/j.jpsychires.2010.10.008)] [Medline: [21035130](https://pubmed.ncbi.nlm.nih.gov/21035130/)]
11. Perednia DA, Brown NA. Tele dermatology: one application of telemedicine. *Bull Med Libr Assoc* 1995 Jan;83(1):42-47 [FREE Full text] [Medline: [7703938](https://pubmed.ncbi.nlm.nih.gov/7703938/)]
12. Coates SJ, Kvedar J, Granstein RD. Tele dermatology: from historical perspective to emerging techniques of the modern era: part I: History, rationale, and current practice. *J Am Acad Dermatol* 2015 Apr;72(4):563-574; quiz 575-576. [doi: [10.1016/j.jaad.2014.07.061](https://doi.org/10.1016/j.jaad.2014.07.061)] [Medline: [25773407](https://pubmed.ncbi.nlm.nih.gov/25773407/)]
13. Kennedy J, Arey S, Hopkins Z, Tejasvi T, Farah R, Secrest AM, et al. Dermatologist perceptions of tele dermatology implementation and future use after COVID-19: Demographics, barriers, and insights. *JAMA Dermatol* 2021 May 01;157(5):595-597 [FREE Full text] [doi: [10.1001/jamadermatol.2021.0195](https://doi.org/10.1001/jamadermatol.2021.0195)] [Medline: [33787839](https://pubmed.ncbi.nlm.nih.gov/33787839/)]

14. Stadler PC, Senner S, Frey S, Clanner-Engelshofen BM, Frommherz LH, French LE, et al. Teledermatology in times of COVID-19. *J Dermatol* 2021 May;48(5):620-624 [FREE Full text] [doi: [10.1111/1346-8138.15812](https://doi.org/10.1111/1346-8138.15812)] [Medline: [33715181](https://pubmed.ncbi.nlm.nih.gov/33715181/)]
15. Kazi R, Evankovich MR, Liu R, Liu A, Moorhead A, Ferris LK, et al. Utilization of Asynchronous and Synchronous Teledermatology in a Large Health Care System During the COVID-19 Pandemic. *Telemed J E Health* 2021;27(7):771-777. [doi: [10.1089/tmj.2020.0299](https://doi.org/10.1089/tmj.2020.0299)] [Medline: [33074786](https://pubmed.ncbi.nlm.nih.gov/33074786/)]
16. Miake-Lye IM, Delevan DM, Ganz DA, Mittman BS, Finley EP. Unpacking organizational readiness for change: an updated systematic review and content analysis of assessments. *BMC Health Serv Res* 2020 Feb 11;20(1):106 [FREE Full text] [doi: [10.1186/s12913-020-4926-z](https://doi.org/10.1186/s12913-020-4926-z)] [Medline: [32046708](https://pubmed.ncbi.nlm.nih.gov/32046708/)]

Abbreviations

APD: Association of Professors of Dermatology

ORCA: Organizational Readiness to Change Assessment

Edited by R Dellavalle, T Sivesind; submitted 25.09.21; peer-reviewed by C Tsou, M Barwick; comments to author 18.03.22; revised version received 04.05.22; accepted 23.05.22; published 09.06.22

Please cite as:

Briggs SM, Lipoff JB, Collier SM

Using Implementation Science to Understand Teledermatology Implementation Early in the COVID-19 Pandemic: Cross-sectional Study

JMIR Dermatol 2022;5(2):e33833

URL: <https://derma.jmir.org/2022/2/e33833>

doi: [10.2196/33833](https://doi.org/10.2196/33833)

PMID: [35720247](https://pubmed.ncbi.nlm.nih.gov/35720247/)

©Shanelle Mariah Briggs, Jules Benjamin Lipoff, Sigrid Marie Collier. Originally published in *JMIR Dermatology* (<http://derma.jmir.org>), 09.06.2022. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in *JMIR Dermatology Research*, is properly cited. The complete bibliographic information, a link to the original publication on <http://derma.jmir.org>, as well as this copyright and license information must be included.